



NUMERICAL REGULARITY MAP FOR FUNDAMENTAL ONE-DIMENSIONAL FRACTIONAL DIFFERENTIAL EQUATIONS WITH HÖLDER CONTINUOUS SOLUTIONS

MANA KATO

Graduate School of Science and Engineering, Doshisha University,
1-3 Tatara Miyakodani, Kyotanabe, Kyoto, 610-0394, Japan
(E-mail: ctwd0905@mail4.doshisha.ac.jp)

HIROSHI FUJIWARA

Graduate School of Informatics, Kyoto University,
Yoshida-honmachi, Sakyo-ku, Kyoto, 606-8501, Japan
(E-mail: fujiwara@acs.i.kyoto-u.ac.jp)

and

HITOSHI IMAI

Faculty of Science and Engineering, Doshisha University,
1-3 Tatara Miyakodani, Kyotanabe, Kyoto, 610-0394, Japan
(E-mail: himai@mail.doshisha.ac.jp)

Abstract. In the paper, fundamental one-dimensional fractional differential equations in the sense of Caputo with Hölder continuous solutions are solved numerically. A large number of numerical experiments results in a numerical regularity map which visualizes relationship between the Hölder exponent of the solution, the convergence rate of schemes, and the order of the derivative. The map reveals that the rate of convergence changes depending on whether the derivative order is greater than or less than one.

Communicated by Editors; Received September 29, 2021

This work was supported by JSPS KAKENHI Grant Numbers 18K03436, 16H02155, 20H01821, 21H00999, and 21K18586.

AMS Subject Classification: 34G20, 34K28, 65L05.

Keywords: fractional derivative, Hölder continuity, spectral collocation method, multiple precision.