



ON THE NECESSARY CONDITIONS FOR PRESERVING THE NONNEGATIVE CONE: DOUBLE SCALE ANOMALOUS DIFFUSION

MESSOUD EFENDIEV *

Helmholtz Zentrum München, Institut für Computational Biology,
Ingolstädter Landstrasse 1 Neuherberg, 85764, Germany

Fen-Edebiyat Fakültesi, Biyoloji Bölümü,
GZFC-212, Göztepe Kampüsü, Kadikoy, Istanbul

(E-mail: messoud.efendiyev@helmholtz-muenchen.de, m.efendiyev@marmara.edu.tr)

and

VITALI VOUGALTER †

Department of Mathematics, University of Toronto
Toronto, Ontario, M5S 2E4, Canada

(E-mail: vitali@math.toronto.edu)

Abstract. The work deals with the easily verifiable necessary conditions of the preservation of the nonnegativity of the solutions of a system of parabolic equations in the case of the double scale anomalous diffusion when the fractional Laplacian is added to the negative Laplace operator raised to another fractional power in the space of two dimensions. Such necessary conditions are extremely important for the applied analysis society because they impose the necessary form of the system of equations that must be studied mathematically.

Communicated by Editors; Received May 2, 3022

This work is partially supported by the Robertson Chair grant (V. V.).

AMS Subject Classification: 35K55, 35K57, 76R50.

Keywords: double scale anomalous diffusion, parabolic systems, nonnegativity of solutions.