



A Note on the Entire Solutions for the System of Partial Differential-Difference Equations of Fermat-Type

JAYASHRI PATTAR

Department of Mathematics,
School of Engineering,
Presidency University, Bengaluru-560064, Karnataka, India.
(E-mail: jspmcu.2019@gmail.com)

and

SHILPA N

Department of Mathematics,
School of Engineering,
Presidency University, Bengaluru-560064, Karnataka, India.
(E-mail: shilpajaikumar@gmail.com)

Abstract. The present study focuses to find out a result for the existence and different forms of entire solutions of Fermat-type for the system of partial differential-difference equations:

$$\begin{cases} \left[\frac{\partial f_1(z_1, z_2)}{\partial z_1} + \frac{\partial f_1(z_1, z_2)}{\partial z_2} \right]^2 + f_2(z_1 + c_1, z_2 + c_2)^2 = e^{g(z_1, z_2)}, \\ \left[\frac{\partial f_2(z_1, z_2)}{\partial z_1} + \frac{\partial f_2(z_1, z_2)}{\partial z_2} \right]^2 + f_1(z_1 + c_1, z_2 + c_2)^2 = e^{g(z_1, z_2)}, \\ \left[\frac{\partial f_1(z_1, z_2)}{\partial z_1} + \frac{\partial f_1(z_1, z_2)}{\partial z_2} \right]^2 + [f_2(z_1 + c_1, z_2 + c_2) - f_1(z_1, z_2)]^2 = e^{g(z_1, z_2)}, \\ \left[\frac{\partial f_2(z_1, z_2)}{\partial z_1} + \frac{\partial f_2(z_1, z_2)}{\partial z_2} \right]^2 + [f_1(z_1 + c_1, z_2 + c_2) - f_2(z_1, z_2)]^2 = e^{g(z_1, z_2)} \end{cases}$$

and

$$\left(\frac{\partial^2 f(z_1, z_2)}{\partial z_1 \partial z_2} \right)^n + f(z_1 + c_1, z_2 + c_2)^m = e^{g(z_1, z_2)}.$$

Our enquiry attempt to generalize and extend the result of H. Y. Xu, et al., [20] to the system of equations.

Communicated by Editors; Received December 11, 2022.

AMS Subject Classification: 30D35, 39A45, 35M10.

Keywords: Entire solution, finite order, linear function, partial differential-difference equations, Fermat-type.