



UNIQUENESS OF L -FUNCTIONS CONCERNING HOMOGENEOUS DIFFERENTIAL POLYNOMIALS WITH WEAKLY WEIGHTED SHARING

HARINA P. WAGHAMORE*

Department of Mathematics, Jnanabharathi Campus, Bangalore University
Bangalore, India - 560 056

(E-mail: harinapw@gmail.com, harina@bub.ernet.in)

and

MEGHA M. MANAKAME

Department of Mathematics, Jnanabharathi Campus, Bangalore University
Bangalore, India - 560 056

(E-mail: megha.manakame80@gmail.com, meghamm@bub.ernet.in)

Abstract. The \mathfrak{L} - functions with Riemann zeta function and its various generalizations have been widely examined by mathematicians worldwide. \mathfrak{L} - functions are analytically continued as meromorphic functions and \mathfrak{L} - functions are Selberg class functions, with the Riemann zeta function serving as the prototype. It is convenient to study the value distribution and uniqueness problems on \mathfrak{L} - functions and arbitrary meromorphic functions. Additionally, we can better examine some of the classical results, since \mathfrak{L} - functions only have a possible pole at $z = 1$. In this paper, by using the concept of weakly weighted sharing, we study the value distribution of a \mathfrak{L} - function and an arbitrary meromorphic function when certain type of homogeneous differential polynomials $\zeta^n Q(\zeta)$ and $\mathfrak{L}^n Q(\mathfrak{L})$ share $\omega(\rho, \iota)$. Our results extends and generalizes some recent results due to X. M. Li et al. (Filomat 33:18, 2019, 5767-5776).

*Corresponding Author

Communicated by Editors,; Received April 13, 2024

AMS Subject Classification: 30D35

Keywords: Meromorphic functions, L - function, Weakly weighted sharing, Uniqueness, Homogeneous differential polynomial etc.