



AN EFFICIENT COMPUTATION OF WAVELETS BASED ON TRIGONOMETRIC B-SPLINES REVERSE SUBDIVISION

AJEDDAR MOHAMED

University Hassan First,
FST, MISI Laboratory, Settat, Morocco.
(E-mail: m.ajeddar@uhp.ac.ma)

and

ABDELLAH LAMNII

University Abdelmalek Essaadi,
ENSATé, ISI Laboratory, Tetouan, Morocco.
(E-mail: a_lamnii@yahoo.fr)

Abstract. This paper presents a novel approach for deriving refinement equations applicable to uniform algebraic trigonometric B-splines (UAT B-splines) of arbitrary order k . In this work, we develop a general formula for the refinement equation tailored to UAT B-splines of any order k . Leveraging the matrix representation of this equation, we further derive the subdivision formula for UAT B-spline curves. Additionally, we provide an explicit form of the inverse subdivision scheme corresponding to quadratic UAT B-splines. The study also delves into constructing mutually orthogonal spline wavelet spaces on uniform partitions of a bounded interval, utilizing reverse subdivision schemes, and substantiates the findings with numerical experiments.

Communicated by Editors; Received September 16, 2024

AMS Subject Classification: 41A15, 65T60, 65D07.

Keywords: Quadratic B-Splines, Refinement Equations, Inverse Subdivision, Multiresolution Analysis, wavelet.