



EVOLUTION QUASI-VARIATIONAL INEQUALITIES OF MAXWELL TYPE

Dedicated to the memory of Professor Mario Primicerio

MARIA GOKIELI

Faculty of Mathematics and Natural Sciences, School of Exact Sciences,
UKSW, Warsaw, Poland
(E-mail: m.gokieli@uksw.edu.pl)

NOBUYUKI KENMOCH

Faculty of Education, Chiba University, Chiba, Japan.
(E-mail: nobuyuki.kenmochi@gmail.com)

and

MAREK NIEZGÓDKA

Silesian University of Technology, Gliwice, Poland
(E-mail: marekn1506@gmail.com)

Abstract. This paper is concerned with evolution quasi-variational inequalities of Maxwell type. Maxwell equation $\mathbf{u}_t + \nabla \times (\rho \operatorname{curl} \mathbf{u}) = \mathbf{f}$ is well known as a fundamental equation, where \mathbf{u} is a magnetic field, ρ is a nonnegative function of resistivity and \mathbf{f} is a given vector field. However in the critical-state (involving superconductivity) the dynamics of \mathbf{u} is quite different from usual situation; for instance the current density $|\operatorname{curl} \mathbf{u}|$ cannot exceed some critical value γ_0 depending on temperature ζ and magnetic field \mathbf{u} itself, like $|\operatorname{curl} \mathbf{u}| \leq \gamma_0(\zeta, \mathbf{u}, \dots)$. Following the suggestion of L. Prigozhin [10], we shall treat a quasi-variational inequality with unknown dependent constraint set $\mathbf{K}(t)$ of the form:

$$\mathbf{u}_t + \partial I_{\mathbf{K}}(\mathbf{u}) + \nabla \times (\rho \operatorname{curl} \mathbf{u}) = \mathbf{f}, \quad \mathbf{K} := \{\mathbf{u} \mid |\operatorname{curl} \mathbf{u}| \leq \gamma_0(\zeta, \mathbf{u}, \dots)\}, \quad (*)$$

in an appropriate function space. We shall precisely reformulate our problem as a system of (*) and a heat equation depending on \mathbf{u} , and show the existence of a weak solution.

Communicated by Editors; Received July 17, 2025

AMS Subject Classification: 34G25, 35K51, 35K57, 35K50.

Keywords: convex, subdifferential, quasi-variational inequality, superconductivity, Maxwell equation